

Charged Black Hole Solutions in Einstein-Born-Infeld gravity with a Cosmological Constant

Sharmanthie Fernando* ¹ and Don Krug** ²

**Department of Physics & Geology*

***Department of Mathematics & Computer Science*

Northern Kentucky University

Highland Heights

Kentucky 41099

U.S.A.

Abstract

We construct black hole solutions to Einstein-Born-Infeld gravity with a cosmological constant. Since an elliptic function appears in the solutions for the metric, we construct horizons numerically. The causal structure of these solutions differ drastically from their counterparts in Einstein-Maxwell gravity with a cosmological constant. The charged de-Sitter black holes can have up to three horizons and the charged anti-de Sitter black hole can have one or two depending on the parameters chosen.

Key words: Born-Infeld, Cosmological constant, Black Holes

1 Introduction

Born-Infeld electrodynamics was first introduced in 1930's to obtain a finite energy density model for the electron [1]. It has attracted considerable interest in recent times due to various reasons. One of the motivations being the observations that it arises naturally in open superstrings and in D-branes [2]. The low energy effective action for an open superstring in loop calculations lead to Born-Infeld type actions [3]. It has also being observed that the Born-Infeld action arises as an effective action governing the dynamics of vector-fields on D-branes [4]. For a review of aspects of Born-Infeld theory in string theory see Gibbons [5].

¹fernando@nku.edu

²krugd@nku.edu

In recent times, the cosmological constant has been considered for several theoretical and observational reasons. One of the strongest supports have been the recent results from a supernova which suggests a positive cosmological constant [6]. From the theoretical point of view, recent advances in string theory have been subjected to studies in all dimensions with a negative cosmological constant [7]. Asymptotically anti-de Sitter black holes are more interesting than their counterparts since they allow more geometries for black hole horizons than for the $\Lambda = 0$ case [8]. Also they are thermodynamically stable which has been an inspiration for much recent work [9]. Therefore studying black holes in (anti)de Sitter spaces has an important place in current research.

In this paper we explore the existence of electrically charged black holes in Einstein-Born-Infeld gravity with a cosmological constant for a U(1) gauge field. Our motivation is to see how the singular nature of black holes gets modified when coupled to non-linear electrodynamics. We find that depending on the value of the cosmological constant Λ , mass parameter M , coupling constant β and the charge Q , the black hole solutions can have up to three horizons.

The paper is organized as follows: In section 2 the basic equations are derived. In section 3 the general solutions to static spherically symmetric solutions are constructed. In section 4 the solutions are studied in detail for various values of the parameters of the theory. In section 5 the temperature is calculated and finally in section 6 the conclusion are given.

2 Basic Equations

In this section we will derive the equations of motion for non-linear electrodynamics. The most general action for such a theory coupled to gravity with a cosmological constant is as follows:

$$S = \int d^4x \sqrt{-g} \left[\frac{(R - 2\Lambda)}{16\pi G} + L(F) \right] \quad (1)$$

Here, $L(F)$ is a function of the field strength $F_{\mu\nu}$ only. For the weak field limit, $L(F)$ has to be of the form

$$L(F) = -F^{\mu\nu} F_{\mu\nu} + O(F^4) \quad (2)$$

In this paper, we will study a particular non-linear electrodynamics called Born-Infeld theory which has attracted lot of attention due to its relation to string effective actions. The function $L(F)$ for Born-Infeld may be expanded to

$$L(F) = 4\beta^2 \left(1 - \sqrt{1 + \frac{F^{\mu\nu} F_{\mu\nu}}{2\beta^2}} \right) \quad (3)$$

Here, β has dimensions $length^{-2}$ and G $length^2$. In the following sections we will take $16\pi G = 1$.

By extremising the Lagrangian in eq.(1), with respect to the metric $g_{\mu\nu}$ and the electrodynamics potential A_μ , one obtains the corresponding field equations as follows:

$$\nabla_\mu \left(\frac{F^{\mu\nu}}{\sqrt{1 + F^2/2\beta^2}} \right) = 0 \quad (4)$$

$$R_{\mu\nu} - g_{\mu\nu}\Lambda = \left(\frac{\partial L(F)}{\partial g^{\mu\nu}} + \frac{g_{\mu\nu}A(F)}{2} \right) \quad (5)$$

where

$$A(F) = -L(F) + \frac{\partial L(F)}{\partial g^{ab}} g^{ab}; \quad \frac{\partial L(F)}{\partial g^{\mu\nu}} = \frac{-2F_{\beta\nu}F_\mu^\beta}{\sqrt{1 + F^2/2\beta^2}} \quad (6)$$

3 Static Spherically Symmetric Solutions

We will study static spherically symmetric charged solutions to Einstein-Born-Infeld gravity with a cosmological constant. The most general metric for such a configuration can be written as

$$ds^2 = -e^{2\mu} dt^2 + e^{2\nu} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) \quad (7)$$

To simplify the equations, we use tetrad formalism and Cartan structure equations. The obvious orthonormal basis is the following:

$$\theta^0 = e^\mu dt; \quad \theta^1 = e^\nu dr; \quad \theta^2 = r d\theta; \quad \theta^3 = r \sin(\theta) d\phi \quad (8)$$

The indices $a, b = 0, 1, 2, 3$ are for the orthonormal basis and $\mu, \nu = 0, 1, 2, 3$, for the coordinate basis with $x^0 = t, x^1 = r, x^2 = \theta, x^3 = \phi$. Now, one can compute the $R_{\mu\nu}$ for the orthonormal basis given in eq.(8) as follows:

$$R_{00} = e^{-2\nu}(\mu'' + \mu'^2 - \mu'\nu' + \frac{2\mu'}{r}) \quad (9)$$

$$R_{11} = e^{-2\nu}(-\mu'' + \nu'^2 + \mu'\nu' + \frac{2\nu'}{r}) \quad (10)$$

$$R_{22} = R_{33} = \frac{1}{r^2}(1 - e^{-2\nu}) + \frac{1}{r}(-\mu' + \nu')e^{-2\nu} \quad (11)$$

For static electrically charged solutions, the only non-zero field strength is F_{rt} or F_{01} . This leads to,

$$\frac{\partial L(F)}{\partial g^{11}} = -\frac{\partial L(F)}{\partial g^{00}}; \quad \frac{\partial L(F)}{\partial g^{22}} = \frac{\partial L(F)}{\partial g^{33}} = 0. \quad (12)$$

Note that the conditions in equation(12) are satisfied by any function L of F^2 for static electrically charged metric. The condition in eq.(12) and the fact that $\eta_{ab} = -, +, +, +$, combined with the above field equations leads to,

$$R_{11} = -R_{00} \quad \Rightarrow \mu + \nu = 0 \quad (13)$$

In order to solve for the function μ and ν , we will use the electromagnetic equations(4). The non-vanishing components of the electrodynamic field tensor in the coordinate basis are given by $F_{tr} = E(r)$ and in the orthonormal basis $F_{01} = \hat{E}$. They are related to each other by, $E(r) = \hat{E}(r)e^{\mu+\nu}$. Hence, the non-linear Lagrangian reduces to,

$$L(F) = 4\beta^2 \left(1 - \sqrt{1 - \frac{E^2}{\beta^2}} \right) \quad (14)$$

Note that the eq.(14) imposes an upper bound for $|E|$ to be smaller than β . This is a crucial characteristic of non-linear electrodynamics which leads to finite self energy of the electron. The electric field E follows from eq.(4) as,

$$E(r) = \frac{Q}{\sqrt{r^4 + Q^2/\beta^2}} \quad (15)$$

From Einstein's equations one can derive a relation for ν as follows,

$$(re^{-2\nu})' = 1 - \Lambda r^2 + 2\beta(r^2\beta - \sqrt{Q^2 + r^4\beta^2}) \quad (16)$$

Hence,

$$e^{-2\nu} = 1 - \frac{2M}{r} - \frac{\Lambda r^2}{3} + 2\beta \left(\frac{r^2\beta}{3} - \frac{1}{r} \int_r^\infty \sqrt{Q^2 + r^4\beta^2} \right) \quad (17)$$

In the limit $\beta \rightarrow \infty$, the elliptic integral can be expanded to give,

$$(re^{-2\nu})' = 1 - \Lambda r^2 - \frac{Q^2}{r^2} \quad (18)$$

resulting the function $\nu(r)$ for the Reissner-Nordstrom-(anti)de Sitter solutions,

$$e^{-2\nu} = 1 - \frac{2M}{r} - \frac{\Lambda r^2}{3} + \frac{Q^2}{r^2} \quad (19)$$

Here M is an integrating constant which may be interpreted as a quasi-local mass when $\Lambda = 0$ [10].

4 Electrically Charged Black Hole Solutions

In this section we will explore the existence of black holes solutions from the above general solutions. For regular horizon to exist the metric function $g_{tt} = e^{-2\nu}$ has to have zeros. First, let us redefine $e^{-2\nu}$ as $f(r)$ and define $g_0(r)$ as the function $d(rf(r))/dr$.

$$f(r) = 1 - \frac{2M}{r} - \frac{\Lambda r^2}{3} + 2\beta \left(\frac{r^2\beta}{3} - \frac{1}{r} \int_r^\infty \sqrt{Q^2 + r^4\beta^2} \right)$$

$$g_0(r) = 1 + (2\beta^2 - \Lambda)r^2 - 2\beta\sqrt{Q^2 + \beta^2 r^4} \quad (20)$$

The number of positive roots of $f(r)$ and of $rf(r)$ are the same. Therefore, the number of roots of $f(r)$ would be at most the number of roots of $g_0(r)$ plus one. (which is a result of Rolle's theorem in calculus). Since we are looking for roots of $g_0(r)$, we will consider the function $g(r)$ which is obtained by simplifying $g_0(r) = 0$.

$$g(r) = r^4(\Lambda^2 - 4\beta^2\Lambda) + r^2(4\beta^2 - 2\Lambda) + (1 - 4\beta^2Q^2) \quad (21)$$

To understand the behavior of the black hole metric for very small r , one can do a series expansion of $f(r)$ around $r = 0$ as follows:

$$f(r) \approx 1 - \frac{(2M - A)}{r} - \frac{10}{3}\beta Q + \frac{2\beta^2}{3}r^2 - \frac{\beta^2}{5}r^4 \quad (22)$$

Here,

$$A = \frac{1}{3}\sqrt{\frac{\beta}{\pi}}Q^{3/2}\Gamma\left(\frac{1}{4}\right)^2 \quad (23)$$

We will define $2M' = 2M - A$ to discuss the behavior of the geometry. Hence when $r \rightarrow 0$, the behavior of $f(r)$ is dominated by $2M'/r$ term irrespective of the value of Q and Λ . One may recall that in the Einstein-Maxwell black hole solutions, it is Q^2/r^2 that dominates the behavior of the space-time around $r = 0$. This is a major difference incorporating Born-Infeld electrodynamics to gravity compared to the Maxwell case.

We will classify the solutions according to the value of the integration constant M' in the following sections. We will consider $\Lambda > 0$ and $\Lambda < 0$ cases separately. Note that the $\Lambda = 0$ case has been studied by Rasheed [11].

4.1 De Sitter Black Holes

For $\Lambda > 0$, $g(r)$ is a 4th degree even function and can have at the most two positive roots. Hence $f(r)$ can have at the most three roots. However, if one takes a closer look at the behavior of the function for $r \rightarrow \infty$ and $r \rightarrow 0$ one can predict the number of roots exactly. From the expansions in eq.(22), it is obvious that for small r , $f(r) \rightarrow -2M'/r$ and for large r , $f(r) \rightarrow -\infty$.

When $M' > 0$, for small r , the function $f(r) \rightarrow -\infty$. Hence $f(r)$ does not have a root at all or it has two roots or it has a degenerate root as shown in the Fig.1. In all these cases, the curvature scalars $R_{\alpha\gamma\beta\rho}R^{\alpha\gamma\beta\rho}$, $R_{\alpha\beta}R^{\alpha\beta}$ and R explode at $r = 0$ and are finite else where. Hence there is a curvature singularity at $r = 0$. Since for large r , $f(r) < 0$, r becomes a time coordinate. Hence the future infinity is space-like and such space-times have cosmological event horizons. Such space-times are similar to Schwarzschild-de Sitter black hole space-time [12].

In the Fig.1, the graph in plain line represents a solution with two roots to the function $f(r)$. The smaller of these values, which can be called r_+ can be regarded

as the position of the black hole event horizon. The larger value of the roots r_{++} represents the position of the cosmological event horizon for observers on the world-lines of constant r between r_+ and r_{++} . The Killing vector $\frac{\partial}{\partial t}$ is null on both these horizons. These horizons can be removed by appropriate coordinate patches similar to the Schwarzschild-de-Sitter black hole space-time [12].

As M' increases, the black hole event horizon increases and the cosmological event horizon decreases. For special values of the M' , the two coincides leading to a degenerate horizon as given in the Fig.1 (dashed).




Figure 1. The figure shows the function $f(r)$ for $\Lambda = 1.5$, $\beta = 0.01$ and $Q = 1$. The lighter graph shows $f(r)$ for $M' = 0.1$ and the dashed one is for $M' = 0.4$.

4.2 Anti de-Sitter Black Holes

For $\Lambda < 0$, one can apply Decarte's rule on the discriminant to predict the behavior of the roots of $g(r)$ as follows; Since $(2\beta^2 - \Lambda) > 0$, $g(r)$ can have only one root leading $f(r)$ to have two roots.

When $M' > 0$ the function $f(r) \rightarrow -\infty$ for small r and goes to ∞ for large r . From the Fig. 2, there is only one horizon. There is a curvature singularity at $r = 0$. Hence this is very similar to Schwarzschild black hole with a space-like singularity at the origin.




Figure 2. The figure shows the function $f(r)$ for $\Lambda = -0.1$, $\beta = 0.05$, $M' = 1$ and $Q = 1$

When $M' = 0$ the function $f(r) \rightarrow (1 - 10\beta Q/3)$ for small r and goes to ∞ for large r . From the Fig. 3, the function $f(r)$ can have one root or none. However, there is no curvature singularity at $r = 0$ and the function $f(r)$ is bounded at $r = 0$. Hence this is a particle-like solution.




Figure 3. The figure shows the function $f(r)$ for $\Lambda = -1$, $M' = 0$ and $Q = 1$. The plain graph is for $\beta = 0.01$, the heavy is for $\beta = 0.5$ and the dashed is for $\beta = 5$

5 Extreme Black Holes

Here we will also discuss the extreme black hole in detail since it is a counterpart to the BPS state in Reissner-Nordstrom solutions. For degenerate roots of $f(r)$, both $f(r)$ and $df(r)/dr$ are zero. Hence from eq.(16), one arrives at the following equation for horizon radii r_{ex} :

$$r_{ex}^4(\Lambda^2 - 4\beta^2\Lambda) + r_{ex}^2(-2\Lambda + 4\beta^2) + (1 - 4Q^2\beta^2) = 0 \quad (24)$$

which gives the solution as,

$$(r_{ex}^2)_{\pm} = \frac{(2\beta^2 - \Lambda) \pm \sqrt{\delta}}{\Lambda(4\beta^2 - \Lambda)} \quad (25)$$

Here,

$$\delta = (2\beta^2 - \Lambda)^2 + (1 - 4\beta^2Q^2)(4\beta^2 - \Lambda)\Lambda \quad (26)$$

The existence of r_{ex} varies for various values of the parameters of the theory as follows:

For $\Lambda < 0$, $(4\beta^2 - \Lambda) > 0$, the denominator of r_{ex} is negative. Since $(2\beta^2 - \Lambda) > 0$, there is at most one positive root for r_{ex}^2 . It occurs only if $(1 - 4\beta^2Q^2)(4\beta^2 - \Lambda)\Lambda > 0$ leading to $\beta Q < 1/2$. The corresponding value is given by r_{ex+}^2 of eq.(24).

For $2\beta^2 > \Lambda > 0$, the denominator of eq.(24) is positive and so is the first element of the numerator. Thus there is always one positive root. A second will occur if $(1 - 4\beta^2Q^2) < 0$ while δ remains positive.

For $4\beta^2 > \Lambda > 2\beta^2$, the denominator of eq.(24) is positive, while the first element of the numerator is negative. There is a positive root if $\delta > 0$ i.e. $\beta Q < 1/2$.

For $\Lambda > 4\beta^2$, both the denominator and the first element of the numerator are negative. So there is always a positive root and there will be two if $(\delta - (2\beta^2 - \Lambda)^2) < 0$ (or $\beta Q > 1/2$) and δ remains positive.

6 Temperature

The Hawking temperature of the black hole solutions discussed above can be calculated as follows: $T = \kappa/2\pi$. Here κ is the surface gravity given by

$$\kappa = -\frac{1}{2} \frac{dg_{tt}}{dr} \Big|_{r=r_+}$$

Here, r_+ is the event horizon of the black hole. Since $f(r) = 0$ at $r = r_+$, the eq.(16) can be used to calculate the surface gravity exactly and the corresponding temperature as,

$$T = \frac{1}{4\pi} \left(\frac{1}{r_+} - \Lambda r_+ + 2\beta \left(r_+ \beta - \frac{\sqrt{(Q^2 + r_+^2 \beta^2)}}{r_+} \right) \right) \quad (27)$$

7 Conclusions

In this paper we presented the local properties of the solutions arising in Einstein-Born-Infeld gravity with a cosmological constant. The non-singular nature of the electric field at the origin changes the structure of the space-times drastically. The singularity at the origin is dominated by the mass term M/r rather than the Q^2/r^2 . Hence the number of horizons changes when compared to the Einstein-Maxwell gravity. Currently we are studying the global properties of these space-times in detail.

In extending this work, we would like to study the stability of these black holes in detail. The electrically charged black holes in anti-de Sitter space have been shown to be unstable for large black holes by using linear perturbation techniques [13]. It would be interesting to study how the non-linear nature effects the instability of such solutions.

From a string theoretical point of view, it is vital to study black hole solutions arising in Einstein-Born-Infeld-dilaton gravity. It is well known that the presence of a dilaton changes the space-time drastically. In such a case, the Lagrangian to be considered would be a more general one with a $SL(2, R)$ invariant Born-Infeld term with a dilaton-axion coupled to it. Solitons and black hole solutions were constructed in a recent paper by Clement and Gal'tsov for Einstein-Born-Infeld-dilaton gravity without a cosmological constant [14]. It would be interesting to study the effects of

both the Born-Infeld term and the dilaton with a cosmological constant which we hope to report elsewhere.

One may recall that the Reissner-Nordstrom AdS black hole is supersymmetric for $Q = 0$ and for $Q^2 = M^2$ which was shown by Romans [15]. It would be interesting to embed the black hole solutions obtained in this paper in a supergravity theory.

Note: The references [16] and [17] was added after the paper was published.

Acknowledgement This work was supported by the grant number: 2001-16 of the Center for Integrative Natural Science and Mathematics of Northern Kentucky University.

References

- [1] M. Born and L. Infeld, Proc. Roy. Soc. Lond. **A144** (1934) 425.
- [2] R. G. Leigh, Mod. Phys. Lett A4 (1989) 2767.
- [3] E. S. Fradkin and A. A. Tseytlin, Phys. Lett. **B163** (1985) 123.
- [4] A. A. Tseytlin, Nucl. Phys. **B276** (1986) 391.
- [5] G. W. Gibbons, hep-th/0106059
- [6] S. M. Carroll, Living Rev.Rel. 4 (2001) 1 , astro-ph/0004075
- [7] O. Aharony, S. S. Gubser, J. Maldesena and H. Ooguri, Phys.Rept. 323 (2000) 183-386.
- [8] L. Vanzo, Phys.Rev. D56 (1997) 6475-6483.
- [9] S. Hawking and D. Page, Comm. Math. Phys. **87** 577 (1983).
- [10] T. Zannias, Phys. Rev. **D41** 3252 (1990).
- [11] D. A. Rasheed, hep-th/9702087.
- [12] G. W. Gibbons and S. W. Hawking, Phys. Rev **D15** 2738, (1977)
- [13] S. S. Gubser and I. Mitra, hep-th/0009126
- [14] G. Clement and D. Gal'tsov, Phys.Rev.**D62** 124013, (2000)
- [15] L. J. Romans, Nucl.Phys. **B 383** 395 (1992).
- [16] D. L. Wilshire, Phys. Rev. **D 638** 2445, (1998)
- [17] G. W. Gibbons and C.A.R. Herdeiro, Class.Quant.Grav.18 1677 (2001)